

Louisiana State University
LSU Digital Commons

Faculty Publications

Department of Oceanography & Coastal
Sciences

2018

Response To The Letter To The Editor On Turner (2017)

R E. Turner
eurne@lsu.edu

Follow this and additional works at: https://digitalcommons.lsu.edu/oceanography_coastal_pubs

Recommended Citation

Turner, R. E. (2018). Response To The Letter To The Editor On Turner (2017). *Geo-Geography And Environment*, 5 (2) <https://doi.org/10.1002/geo2.62>

This Article is brought to you for free and open access by the Department of Oceanography & Coastal Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

COMMENTARY

WILEY



Response to the letter to the editor on Turner (2017)

R. Eugene Turner 

Department of Oceanography and Coastal Sciences, Louisiana State University, Baton Rouge, LA, USA

Correspondence

Robert Eugene Turner

Email: euturne@lsu.edu

This commentary responds to Schueller et al. (2018), <https://doi.org/10.1002/geo2.48>.

The authors of the letter to the editor (LTE; Schueller et al., 2018) raise three objections to the analysis of Turner (2017): (1) using an estimate of MSY (maximum sustainable yield) as a stock management metric; (2) inappropriate analyses; and (3) failure to consider “alternative hypotheses for the putative patterns.”

1. They write “neither value described as ‘MSY’ in Turner (2017) represents an accurate or reliable estimate of MSY for menhaden.” The MSY value used is defined by them as a “biological reference point.” That is why it was used; it is the maximum equilibrium landing “against which to measure stock or fishery status” and includes a spawner–recruit curve and values for growth, mortality and gear selectivity (Williams & Shertzer, 2015). The MSY usually serves as “a value to be avoided” (Vaughan et al., 2007). The MSY values are indeed, as the LTE author(s) write, “reference points for management.” If the LTE authors mean that the estimated MSY in SEDAR (2013) is incorrectly estimated by the NMSF Beaufort Assessment Model (BAM), then that concern should be addressed to the 27 contributors to the BAM development, and to the several reviewers involved in the two lengthy review cycles. The status today is that the MSY used is an authoritative benchmark metric developed using output from the BAM.
2. The analysis of data used in their Table 1 incorrectly attributes similarity to the data used in Turner (2017), which come from table 4 in Vaughan et al. (2010) and are repeated in tables 3.5 and 3.6 of SEDAR (2013). The data are not “mean estimates of size-at-age,” as categorised in the LTE, but weighted averages. For example, the table heading for the data includes: “Weighted mean fork length (mm) at age, with weightings based on annual catch in numbers by season and area” and “Weighted mean weight (g) at age, with weightings based on annual catch in numbers by season and area.” These data are, therefore, the result of accounting for different numbers of fish taken from different areas and in different months. The LTE authors used the raw data for *all* fish regardless of month, and without discrimination for the number of fish each month, growth between months, or catch area – a crude approach to calculating average size that obviously adds variation to the “average” value for each year; it is not the data set used in Turner (2017) and the comment is, therefore, an inappropriate comparison.
3. The first sentence of the abstract (Turner, 2017) states the hypothesis tested: “to determine if there was evidence of changes consistent with the well documented temperature size rules.” The decline in size per degree (slope) is the same when fishing is below the MSY compared with when it is above the MSY (Figure 1), but there is a smaller size with greater fishing intensity for each age group. Support for a temperature effect is, therefore, demonstrated as being consistent with the hypothesis tested and influenced by fishing intensity.

Did the size change with temperature, in general? Yes. Did the analysis show distinction between when fish were over and under a standard metric of fishing intensity? Yes. Perhaps there is another explanation for the trends observed; but the most likely one today is consistent with results from a broad suite of analyses about the effect of warming oceans on organisms

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

The information, practices and views in this article are those of the author(s) and do not necessarily reflect the opinion of the Royal Geographical Society (with IBG).

© 2018 The Author. *Geo: Geography and Environment* published by John Wiley & Sons Ltd and the Royal Geographical Society (with the Institute of British Geographers).




FIGURE 1 Figure 4 in Turner (2017) is sub-divided here into four panels for emphasis. The data are the average weight of fish for each age class of Gulf of Mexico menhaden. The dataset is further divided into years when the total harvest was less than or greater than the modelled maximum sustainable yield (MSY). The slopes were not significant among year classes, whereas the intercept was different.

(Baudron et al., 2014; Daufresne et al., 2009; Ohlberger, 2013; Pauly & Cheung, 2017; Sheridan & Bickford, 2011). It is inconsistent with other hypotheses mentioned in the paper.

The commenters correctly point out that the Y axis in Figure 1 (Turner, 2017) should be “thousand mt,” not “million mt.” I regret the error, and it does not change the results.¹

ACKNOWLEDGEMENTS

This research was made possible by a grant from The Gulf of Mexico Research Initiative. Data are publicly available through the Gulf of Mexico Research Initiative Information & Data Cooperative (GRIIDC) at <https://data.gulfresearchinitiative.org> (<https://doi.org/10.7266/n77d2s57>).

ENDNOTE

¹ This error was corrected online on 11 September 2018.

ORCID

R. Eugene Turner  <http://orcid.org/0000-0003-0776-7506>

REFERENCES

- Baudron, A. R., Needle, C. L., Rijnsdorp, A. D., & Marshall, C. T. (2014). Warming temperatures and smaller body sizes: Synchronous changes in growth of North Sea fishes. *Fisheries Bulletin*, 20, 1023–1031. <https://doi.org/10.1111/gcb.12514>
- Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in aquatic ecosystems. *Proceedings National Academy of Sciences of the United States of America*, 106, 12788–12793. <https://doi.org/10.1073/pnas.0902080106>
- Ohlberger, J. (2013). Climate warming and ectotherm body size from individual physiology to community ecology. *Functional Biology*, 27, 991–1001. <https://doi.org/10.1111/1365-2435.12098>

- Pauly, D., & Cheung, W. L. (2017). Sound physiological knowledge and principles in modeling shrinking of fishes under climate change. *Global Change Biology*, 23, 1–12. <https://doi.org/10.1111/gcb.13831>
- Schueller, A. M., Leaf, R. T., Mroch III, R. M., & Nessler, G. M. (2018). Response to Turner. *Geo: Geography and Environment*, 5, e00048. <https://doi.org/10.1002/geo2.48>
- SEDAR (2013). *SEDAR 32A – Gulf of Mexico menhaden stock assessment report*. North Charleston, SC: SEDAR.
- Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, 1, 401–506.
- Turner, R. E. (2017). Smaller size-at-age menhaden with coastal warming and fishing intensity. *Geo: Geography and Environment*, 4, e00044. <https://doi.org/10.1002/geo2.44>
- Vaughan, D. S., Shertzer, K. W., & Smith, J. W. (2007). Gulf menhaden (*Brevoortia patronus*) in the U.S. Gulf of Mexico: fishery characteristics and biological reference points for management. *Fisheries Research*, 83, 263–275. <https://doi.org/10.1016/j.fishres.2006.10.002>
- Vaughan, D. S., Smith, J. W., & Shertzer, K. W. (2010). Population Characteristics of Gulf Menhaden. *Brevoortia patronus NOAA Technical Report NMFS 149 A Technical Report of the Fishery Bulletin*.
- Williams, E. H., & Shertzer, K. W. (2015). *Technical documentation of the Beaufort Assessment Model (BAM)*. US Department of Commerce, NOAA Technical Memorandum NMFS–SEFSC–671.

How to cite this article: Turner RE. Response to the letter to the editor on Turner (2017). *Geo: Geography and Environment*. 2018;e00062. <https://doi.org/10.1002/geo2.62>